

THE EVOLUTIONARY PHYSIOLOGY OF ANIMAL FLIGHT: Paleobiological and Present Perspectives

Robert Dudley

*Section of Integrative Biology, University of Texas, Austin, Texas, 78712 and
Smithsonian Tropical Research Institute, P.O. Box 2072, Balboa, Republic of Panama;
e-mail: r_dudley@utxvms.cc.utexas.edu*

Key Words evolution, gigantism, hyperoxia, insect, oxygen

■ **Abstract** Recent geophysical analyses suggest the presence of a late Paleozoic oxygen pulse beginning in the late Devonian and continuing through to the late Carboniferous. During this period, plant terrestrialization and global carbon deposition resulted in a dramatic increase in atmospheric oxygen levels, ultimately yielding concentrations potentially as high as 35% relative to the contemporary value of 21%. Such hyperoxia of the late Paleozoic atmosphere may have physiologically facilitated the initial evolution of insect flight metabolism. Widespread gigantism in late Paleozoic insects and other arthropods is also consistent with enhanced oxygen flux within diffusion-limited tracheal systems. Because total atmospheric pressure increases with increased oxygen partial pressure, concurrently hyperdense conditions would have augmented aerodynamic force production in early forms of flying insects. By the late Permian, evolution of decompositional microbial and fungal communities, together with disequilibrium in rates of carbon deposition, gradually reduced oxygen concentrations to values possibly as low as 15%. The disappearance of giant insects by the end of the Permian is consistent with extinction of these taxa for reasons of asphyxiation on a geological time scale. As with winged insects, the multiple historical origins of vertebrate flight in the late Jurassic and Cretaceous correlate temporally with periods of elevated atmospheric oxygen. Much discussion of flight performance in *Archaeopteryx* assumes a contemporary atmospheric composition. Elevated oxygen levels in the mid- to late Mesozoic would, however, have facilitated aerodynamic force production and enhanced muscle power output for ancestral birds, as well as for precursors to bats and pterosaurs.

INTRODUCTION

Today's oxidizing atmosphere, with an oxygen concentration of about 21%, derives in part from the metabolic activity of photosynthetic organisms. Starting with the anoxic conditions and reducing atmosphere of the Archean and early Proterozoic eons, evolution of cyanobacteria, sulfide-oxidizing bacteria, and algae contributed to buildup of oxygen partial pressures by the beginning of the Cam-

brian (570 MYa) to values likely comparable to those of the present day (1–6). Increased atmospheric oxygen, in turn, may have been an important factor underlying radiations of metazoan taxa in the late Proterozoic and at the base of the Cambrian (7–14). The paleobiological significance of Proterozoic oxygen levels for animal evolution in aquatic habitats has long been recognized, as has been the role of gradual oxygen buildup for subsequent evolution of plant and animal terrestrial communities (6, 8, 9). Less well-studied, however, are potential fluctuations in atmospheric constituents through the course of the Phanerozoic.

In particular, variation in oxygen partial pressure and concomitant variation in air density potentially influence the physiology and biomechanics, respectively, of animal flight performance. Geophysical analyses suggest the presence of a late Paleozoic oxygen pulse beginning in the late Devonian and continuing through to the late Carboniferous. Oxygen concentrations ranged possibly as high as 35% during the initial period of insect flight evolution (late Devonian or early Carboniferous). Atmospheric hyperoxia may thus have physiologically facilitated the initial evolution of insect flight metabolism. Because total atmospheric pressure increases with an increased oxygen partial pressure, concurrently hyperdense conditions would have augmented aerodynamic force production in ancestral winged insects. The multiple historical origins of vertebrate flight similarly appear to correlate temporally with periods of hyperoxia. In particular, elevated oxygen levels in the mid- to late Mesozoic may have facilitated aerodynamic force production and enhanced muscle metabolism for ancestral birds, as well as for precursors to bats and pterosaurs. Because of such implications for the evolution of animal flight performance, discussion of the mechanisms underlying substantial variation in composition of the earth's atmosphere is warranted.

PHANEROZOIC VARIATION IN THE EARTH'S ATMOSPHERE

The physical properties of the earth's atmosphere are determined almost exclusively by two major constituents, namely diatomic oxygen and diatomic nitrogen. Nitrogen content of the atmosphere is thought to have been approximately constant through geological time (15–17), although some variation in this constituent may derive through biotic linkages with the atmospheric pool (CD Nevison, unpublished data). Various models for oxygen content of the atmosphere in Phanerozoic times have suggested major fluctuations that are driven biotically (6, 8, 13, 16, 19, 20). In particular, geochemical modeling (21, 22) suggests a late Paleozoic disequilibrium between the rate of carbon fixation by terrestrial plants and the rate at which this fixed material was decomposed and recycled. Through the mid- to end-Devonian (390–362 MYa), this disequilibrium led to a moderate increase in oxygen concentration from 18 to 20%, followed by a more substantial

rise by the end of the Carboniferous (290 MYa) to remarkable values as high as 35% (see Figure 1). Thereafter, an equally pronounced decline resulted in end-Permian and early Triassic values as low as 15%. Relative to the present atmospheric level of 21% (PAL), such hyperoxic and subsequently hypoxic conditions must have had major biological effects during the Paleozoic (23–26). Moreover, atmospheric hyperoxia may also have characterized portions of the Mesozoic. According to the aforementioned geochemical model (21), a secondary rise in oxygen levels characterized the the mid-Jurassic and the Cretaceous (170–65 MYa), with oxygen concentration increasing to 25 to 30% and then declining to PAL only by the end-Tertiary (Figure 1). Such postulated changes in oxygen concentration are at sharp odds with any uniformitarian perspective of atmospheric composition.

The various assumptions underlying reconstruction of ancient atmospheric composition obviously require examination. Current models for Phanerozoic oxygen and carbon dioxide concentrations (21, 27–29) are based on estimates of the exchange rates of fixed and reduced carbon among atmospheric, oceanic, and sedimentary reservoirs. Variation in rates of carbon exchange is driven primarily by the effects of plant terrestrialization, variable rates of organic carbon deposition and decomposition, and changes in continental weathering (20, 22, 30–32). An additional feedback loop potentially influencing atmospheric oxygen levels involves marine phosphorus (33, 34). Because the relative importance of these different mechanisms can be, in many cases, only broadly constrained, considerable uncertainty must be associated with such efforts that model global geochemical cycles. Of particular relevance to the late Paleozoic atmosphere,




Figure 1 Estimate of atmospheric oxygen concentrations through the Phanerozoic (21). PAL: present atmospheric level (20.95%).

however, are the changes necessarily associated with the global phenomenon of terrestrialization by plants. During this process, photosynthetic production of oxygen became decoupled from breakdown of fixed carbon. An ensuing compositional bottleneck, due apparently to the absence of biotic processes leading to rapid breakdown, resulted in extensive deposition of coal and other carbonates. Subsequent evolution by the mid-Permian of microbial decomposers, in particular fungi, then depleted atmospheric oxygen via decay processes and inhibited further carbon accumulation (20, 31, 35). Biological underpinnings to a late Paleozoic oxygen pulse are thus well-grounded, although more precise estimates of excursions in atmospheric carbon dioxide and oxygen await further investigation.

Constraints on fire processes can be used to evaluate the validity of geophysical reconstructions of ancient atmospheres. The models of atmospheric oxygen considered here suggest concentrations approaching but not exceeding 35%, a value that represents an approximate threshold for sustained combustion of the terrestrial biosphere (36, 37). Proliferation of terrestrial ecosystems would clearly be precluded under such circumstances, and analysis of charcoal deposits and of the vegetational fossil record demonstrates that atmospheric oxygen concentrations have not exceeded this upper bound (38). At the lower extreme, the minimum oxygen concentration required for contemporary natural fuels to ignite is about 13% (36, 39); the regular presence of charcoal in the fossil record suggests that atmospheric oxygen never fell below this level (40–42). Methodologically, it is important to realize that ignition of homogeneous woody or paper materials in modern laboratory contexts is not functionally equivalent to the propagation of fire in natural ecosystems. Extrapolation of experimental results with contemporary fuels to the prediction of fire regimes in Paleozoic vegetation is necessarily imprecise, although some inference is possible (34). In sum, these considerations suggest that atmospheric oxygen levels have been historically constrained between 13 and 35%; the best estimate derived from atmospheric modeling (15–35%; Reference 21) is nominally congruent with this range.

No direct indicator is available to test empirically these predictions for ancient levels of atmospheric oxygen. However, diverse geochemical and isotopic evidence is consistent with substantial late Permian declines in both marine and atmospheric oxygen concentrations, declines that likely contributed to the end-Permian extinction event (43–51). Furthermore, atmospheric models incorporating effects of plant terrestrialization suggest substantial changes in the concentration of carbon dioxide. Both modeling and empirical observations demonstrate an approximate 10-fold reduction in carbon dioxide from the mid- to late Paleozoic (22, 27–29, 53–56). For plants, the physiological effects of this draw-down in carbon dioxide are evident in paleontological analyses of stomatal density and other proxies of photosynthetic activity (56–58). Shortcomings of atmospheric modeling notwithstanding, a diversity of empirical evidence is consistent both with carbon dioxide drawdown and with an oxygen pulse in the mid- to late Paleozoic, and with enhanced levels of oxygen in the mid- to late Mesozoic and in much of the Cenozoic.

THE ORIGINS OF ANIMAL FLIGHT

Powered flapping flight evolved independently four times during the Phanerozoic, once in the insects and three times among vertebrates. Evolutionary origins of all volant taxa remain controversial, but both insects and the three taxa of flying vertebrates may have originated during periods of atmospheric hyperoxia. Flight is both biomechanically and physiologically demanding, and the evolution of flight requires dramatic up-regulation of metabolic capacity in addition to the expression of flapping structures that generate useful aerodynamic forces. Although wings are not homologous between insects and volant vertebrates, common physical mechanisms of flight suggest that historically elevated concentrations of atmospheric oxygen have yielded similar aerodynamic consequences during flight evolution.

Wings and Aerodynamic Force Production

Fundamental to force generation by flapping animal wings is the physical nature of the medium to which the wings transfer momentum (59). Air density, the mass of air per unit volume, is a principal determinant of force production by airfoils, biological and otherwise; aerodynamic forces on structures tend to increase linearly with air density. Also, air responds to applied forces by resisting not deformation but rather the rate of deformation, as determined by the viscosity or the internal resistance to flow. For an object such as a wing moving in a fluid, the ensuing reaction force acting parallel to the direction of movement is termed drag. Presence of the wing removes momentum and thus kinetic energy from the moving fluid system, and acts to slow the wing relative to flow. This pressure drag, also known as inertial drag, derives from disruption of the inertial characteristics of the moving flow field and varies with fluid density, relative fluid velocity, and the wing's shape.

Viscous drag, by contrast, emerges from boundary interactions between the wing and the surrounding fluid. Viscosity ensures resistance to flow between adjacent layers of fluid, engendering shear forces within the boundary layer over the wing and imposing a reaction force throughout the fluid that terminates at the wing's surface. Viscous drag thus varies not with density but with fluid viscosity, the relative fluid velocity, and the total wetted surface area of the wing. Pressure and viscous drag have in common a dependence on relative fluid velocity and on object dimensions, but differ fundamentally in their dependence on fluid density and viscosity, respectively. The ratio of inertial to viscous forces on an object moving within a fluid will therefore vary with the ratio of density and viscosity, and must additionally change with object dimensions and its relative velocity. The simplest dimensionless formulation of these four parameters is the Reynolds number (Re), which is proportional to the ratio of inertial to viscous forces that act on objects moving within fluids. Different situations of fluid flow are physically equivalent if the corresponding Re s are approximately the same. In contrast,

variation in either the density or viscosity of a fluid alters the relative magnitude of inertial and viscous forces.

Just as drag is the force parallel to flow around an object, lift is defined as the component of force orthogonal to flow and thus perpendicular to drag in a two-dimensional perspective. For wings at low angles of attack, air flows smoothly over both dorsal and ventral wing surfaces. Because pressure drag of well-designed airfoils is low, momentum extraction from the flow field is minimized. The positive camber of the wing and the downward deflection of air near the trailing wing edge yield slightly different translational velocities for the dorsal and ventral airstreams—airflow is slightly faster above and slower beneath the airfoil. Bernoulli's Principle indicates that this difference in dorsal and ventral airstream velocities results in a pressure gradient; a net force (i.e. lift) acts dorsally upon the wing. In a cross-sectional perspective, air appears to move anteriorly from the ventral to dorsal wing surface and to yield a net rotational movement of air around the wing. This motion of air is equivalent to a rotating flow field or vortex that circulates around the wing and that is centered about the wing itself. For every bound wing vortex, a starting vortex of comparable magnitude but of opposite sense is shed into the wake, ensuring conservation of angular momentum in the fluid.

The physical origin of lift thus lies within the creation of net air circulation about a translating wing. As a wing translates in space, the pressure gradient underlying lift production yields airflow not only around spanwise wing sections (the bound circulation), but also around the wing tip, creating the so-called tip vortex. The tip vortex is linked to the starting and the bound wing vortices and creates a closed vortex loop that exerts a momentum flux on the surrounding air. The downward momentum induced by the presence of the vortex loop is accordingly proportional to the mass of the air moved ventrally (i.e. to the air density) and to the velocity at which the air moves (i.e. the circulation). Lift production thus increases with increased air density (59). The Re at which a wing is operating also exerts a strong influence on lift production. Under highly viscous circumstances, vortex generation becomes more difficult as intermolecular stickiness progressively impedes rotational motions of airflow. Circulatory lift becomes increasingly more difficult to maintain at low Re , whereas the effects of viscous drag also become more pronounced. The lift:drag ratio must then decrease at lower Re , as confirmed empirically for insect and vertebrate wings (60, 61).

Lifting characteristics of wings are thus strongly dependent on both air density and the Re at which they operate. Concomitant with large-scale changes in oxygen concentration during the Phanerozoic, numerous physical characteristics of air must vary if, as suggested previously, nitrogen content of the atmosphere remains constant (23). Parameters such as heat capacity of air, thermal conductivity, diffusivity, and the speed of sound would all have substantially changed under hyperoxic conditions. Of particular interest for the evolution of animal flight is the possibility of density-mediated increases in aerodynamic performance during initial periods of wing evolution (23, 25). The predicted value of air density at the

peak of the late Paleozoic oxygen pulse (285 MYa) is about 1.56 kg/m^3 , an increase of 29% relative to the present sea-level value of 1.21 kg/m^3 . The relative changes in the viscosity of air are, by contrast, much smaller (23). Winglets or proto-wings moving in hyperoxic atmospheres would thus experience a higher air density and would be moving at a comparably increased Re . Both conditions would have facilitated lift production in early flying forms, although the magnitude of these aerodynamic effects would have varied both with morphological features and with the particular wing and body motions of the taxon concerned. Because insect and vertebrate wings are so distinct both ontogenetically and anatomically, origins of flight in the two groups are considered separately.

Origins of Flight in Insects

The evolution of insect flight, a major event in biotic history, can be correlationally linked to atmospheric hyperoxia. The origins of winged (pterygote) insects are indeterminate but probably lie in the Upper Devonian or early Lower Carboniferous. Wingless hexapods are known from 395–390 MYa (62–65), whereas fossils of pterygote hexapods (i.e. winged insects) date from approximately 325 MYa (66, 67). By the Upper Carboniferous, pterygotes are impressively diversified into about fifteen orders (68–71). Although pterygote insects are likely monophyletic (72–78), the morphological origins of wings remain obscure. Wings have been proposed to derive either from fixed paranotal outgrowths of thoracic and abdominal segments in terrestrial taxa (79–86), or from ancestrally mobile gills, gill covers, leg structures, or styli in aquatic forms (87–95). Unfortunately, no transitional forms are known between the wingless apterygotes and the winged pterygote insects, and the biology of early protopterygote forms remains speculative and contentious. Of particular interest to the origins of flight is ancestral habitat association of protopterygotes—were these animals terrestrial or aquatic?

Most evidence, particularly that relating to the physiology and origins of the insect tracheal system, indicates that winged insects are derived from terrestrial apterygote ancestors (61, 96–99). Aquatic larvae, particularly those of the extant and phylogenetically basal Odonata and Ephemeroptera, appear to be secondarily derived (73, 99, 100). Independent of habitat association, however, both larvae and adults of ancestral winged insects probably expressed lateral lobed structures on the abdominal as well as the thoracic segments (68, 90, 101–103). If winglets or wings derived initially from fixed paranotal lobes, flapping motions might have emerged indirectly through action of dorsoventral leg muscles that insert on the thorax, as characterizes so-called bifunctional muscles in many extant insects (104–108). A general question relating to wing origins concerns the possible evolution of novel wing-like structures, as opposed to modification of pre-existing morphological features (85). Acquisition of wings from ancestrally mobile structures might seem more parsimonious than the derivation of flapping wings from stationary paranotal lobes, although the neontological and paleontological data

available at present are insufficient to decide unequivocally between these two hypotheses (61).

A variety of possible functional roles have been attributed to winglets or wings of protopterygotes, including aerodynamic utility, epigamic display during courtship, and thermoregulation (109–115). Hydrodynamic use for what ultimately became aerodynamic structures has been proposed for ancestrally aquatic protopterygotes, as have been possibly amphibious lifestyles (116, 117). Protopterygotes could also have used wing-like structures in air either to drift passively or to skim actively along water surfaces, as do many extant insect taxa (118–122). This behavior is probably a derived rather than a retained ancestral trait of winged insects (61, 123, 124). Although improbable for reasons outlined above, biomechanical considerations suggest that aquatic protopterygotes would have been unlikely to evolve wings that served aerodynamic functions. Water and air differ by almost three orders of magnitude in density, with a corresponding difference in the Re and in the nature of forces generated by oscillating structures. The functionality of wing designs intermediate to either hydrodynamic or aerodynamic force generation is correspondingly unclear (61). Forces of surface tension would present a formidable physical barrier to partial body emergence as well as to projection and oscillation of flattened structures, particularly for the body sizes (2–4 cm) likely characteristic of ancestral pterygotes (64, 125, 126).

Given the assumption of terrestrial pterygote ancestors, a standard explanation for the evolution of wings has been that these structures aerodynamically facilitate jumping escapes from predators on land. Suggestively, a suite of morphological and behavioral protoadaptations for jump-mediated glides are evident among extant apterygote hexapods, the terrestrial sister taxon of the winged insects. Thoracic paranotal lobes as well as styli on the legs and abdominal segments of extant apterygotes could potentially have served in ancestral taxa to generate lift and to facilitate saltatorial escape (61). Neurobiological studies also support the ancestral presence of dedicated sensorimotor pathways underlying escape behavior in both apterygotes and pterygotes (127–129). The startle response of ancestral apterygote insects was then apparently co-opted during pterygote evolution to stimulate jumping, wing flapping, and even evasive flight once airborne (130–137). The historical context of early pterygote evolution was appropriate for imposition of intense predatory pressure by both invertebrates and vertebrates, with a diversity of insectivorous arthropods (particularly arachnids), amphibians, and reptiles found in Devonian and Carboniferous terrestrial ecosystems (138–142). Various morphological characteristics evident among the early Upper Carboniferous entomofauna are also consistent with the hypothesis of predatory defense (141, 143, 144).

Furthermore, the increasing arborescence and geometrical complexity of terrestrial vegetation through the Devonian and into the Carboniferous would have provided a three-dimensional substrate suitable for jumping escapes and ultimately the evolution of flight in protopterygotes (145). Terrestrial vegetation during the period of pterygote origins was composed of fern-like lycopsids and

psilopsids, together with arborescent lycopods, sphenopsids, and progymnosperms (146–150). Lateral jumps from branches or leaves, and particularly jumps directed downward, would have yielded the increased translational velocities necessary for substantial buildup of lift on both thoracic and abdominal projections. Access to nutritional resources, appropriate microhabitats, and suitable oviposition sites would have been facilitated by such an increased capacity for three-dimensional movements. Increased feeding on plants would have been one outcome of enhanced mobility. For example, non-insect hexapods (e.g. *Collembola*) that feed on living plant tissue are rare, whereas approximately 85% of extant insect species are phytophagous at some stage in their life cycle (151). Most tellingly, the fossil record confirms feeding on plants by insects in the Upper Carboniferous and early Permian (152–154).

Given this paleobiological background to wing evolution in insects, it is instructive to consider two distinct features of atmospheric hyperoxia that may have contributed to the evolution of flight. Biomechanically, the greater density of a hyperoxic atmosphere would yield enhanced aerodynamic characteristics of both winglets and the bodies of protopterygotes. Alternatively phrased, less winglet area would be necessary to generate comparable forces if air density were higher, although the magnitude of this effect is strongly dependent on the nature of vortex production around the body and winglets and on the relative air velocity and/or acceleration during takeoffs. Aerodynamic forces on both the body and winglets of protopterygotes would enhance aerial escape during either jumping or steady-state glides (115, 125, 155). The de facto increase in Re associated with increased air density may furthermore have contributed to greater lift production by winglets, particularly for the low aspect ratio wings and Re relevant to pterygote evolution (61, 156). In contrast to theories of wing evolution that require aquatic protopterygotes and a discontinuous selection regime across the air-water interface, escape jumping in terrestrial protopterygotes would from the outset favor gradual and continuous improvement in lift generation. Winglet mobility would then enhance force production, and additionally could be used to control body orientation during flight and while landing (109, 125, 157, 158). Increased air density and higher Re would have been advantageous in any evolutionary scenario involving lift production by winglets or wings. Greater air densities may also have facilitated flight in progressively larger taxa, leading to the giant pterygotes of the late Paleozoic, as discussed below (61, 159).

Physiologically, an increased oxygen content of the late Paleozoic atmosphere would have favored higher and ever more sustained levels of the oxidative metabolism required for flight. As wing flapping became more rapid and of greater amplitude, the energetic costs of flight would have increased substantially; thoracic muscles of extant insects in flight exhibit the highest mass-specific rates of oxygen consumption known for any locomotor tissue (61, 160, 161). This demand for oxygen is met by the air-filled tracheal system, a branching network of cuticular tubes that functions primarily through gaseous diffusion rather than by convective air movement (162–165). Both diffusion and tracheal convection

represent potentially rate-limiting steps in insect respiration, although the much reduced thoracic diameters of small insects (e.g. *Drosophila*) probably alleviate any metabolically based need for active ventilation of the tracheal system. In larger insects, however, diffusional limits on oxygen supply likely constrain maximum body size. For example, flight metabolic rates in an extant dragonfly species vary directly with ambient oxygen concentration (166), a result consistent with diffusion-limited oxygen delivery by the tracheal system. In the late Paleozoic, diffusion of oxygen to the respiring flight muscles would obviously have been enhanced by greater partial pressures of this gas, as well as by the increased diffusion constants associated with an increased total pressure of the atmosphere (25).

Moreover, a spectacular bioindicator of atmospheric hyperoxia is provided by gigantism among late Paleozoic terrestrial arthropods (23, 25, 26). Gigantism in the Carboniferous and Permian was characteristic of many arthropod taxa, including a number of phylogenetically unrelated pterygote orders (141, 167, 168) as well as apterygote insects (169) and other arthropods, including diplopods, arthropleurids, and arachnids (138, 141, 170–172). Although a number of non-mutually exclusive selective forces may have contributed to the evolution of gigantism (90, 141, 173), relaxation of diffusional constraints on maximum body size through atmospheric hyperoxia seems to be the most parsimonious explanation for this phenomenon (19, 23, 25, 174, 175). Necessarily global effects of hyperoxia on animal physiology were thus manifested in the form of dramatically increased body size in diverse arthropod lineages. Furthermore, all giant arthropod taxa of the late Paleozoic went extinct by mid- to end-Permian (23, 25, 168), as would be predicted by the concomitant decline in atmospheric oxygen concentration. Also consistent with the hypothesis linking atmospheric oxygen and diffusional constraints on insect body size is a secondary peak of gigantism (e.g. among mayflies) in the Cretaceous at a time of elevated oxygen concentrations (25, 168). Systematic analysis of Paleozoic gigantism has been precluded to date because of generally poor fossil preservation and incomplete knowledge of interspecific body size distributions. However, detailed analysis of phyletic size change within diffusion-limited taxa does represent an important bioassay for possible effects of atmospheric hyperoxia in the late Paleozoic and the Cretaceous/Tertiary.

Origins of Vertebrate Flight

As with winged insects, the historical origins of flying vertebrates are known only imprecisely. Nonetheless, circumstantial evidence is at least consistent with origins of bats, birds, and pterosaurs during times of hyperdense and hyperoxic atmospheres. The most recent of volant vertebrates are the bats, with a modern morphology apparent in a microchiropteran fossil from 50 MYa. Given this date, chiropteran origins would appear to lie within the early Paleocene or late Cretaceous (176, 177). A similar appearance during hyperoxic times seems likely for birds in the mid- to late Jurassic (178–180). The timing of pterosaur origination

and early diversification is unknown but possibly lies within the Permian given well-developed pterosaur morphologies by the mid- to late Triassic (181, 182). Because a detailed fossil record is unavailable for any of these three taxa, biomechanical analysis of the transitional forms of flight is seriously constrained. Independent of the behavioral or ecological context of flight, however, greater atmospheric density would facilitate aerodynamic force production, whereas increased oxygen partial pressures would similarly enhance oxygen transport to and within the muscles powering flight. Aforementioned discussions of vertebrate flight origins generally assume atmospheric composition of the present day, although the implications of flight evolution in physically variable atmospheres clearly deserve further attention. In a similar vein, reconstructions of ancestral flight physiology that are based on comparisons with extant taxa (183–185) may not fully reflect aerial performance of now extinct taxa in hyperoxic and hyperdense atmospheres.

Most scenarios of vertebrate flight evolution presume jumping takeoffs (directed either with or against gravity), whereas jump-initiated gliding origins of flight are biomechanically more parsimonious than “ground-up” hypotheses for both volant vertebrates and insects (60, 61, 186–188). Jumping via a startle response is widespread among animals (189), and one potential commonality among volant vertebrates and pterygote insects is acquisition of active flight via the pathway of jumping and subsequent gliding to escape predation. Also, unsteady aerodynamic performance during the accelerating portion of a jump may be of greater relevance to escape success than steady-state glides (61). Jump-initiated glides that increased survivorship during predation attempts would potentially select for greater aerodynamic performance, an effect likely to be enhanced at the higher Re of hyperdense air. That predation is a major factor underlying the evolution of aerial locomotion is suggested by the increased longevity of flying animals relative to their non-volant counterparts. For example, volant endotherms have mortality rates significantly lower than those for non-volant endotherms of equivalent body mass (190–193). Paleophysiological simulations of jumping performance can be carried out in extant gliding and flying taxa using variable-density gas mixtures (194), and potentially can be used to investigate the ancestral role of jumping and takeoff aerodynamics in flight evolution.

CONCLUSIONS AND IMPLICATIONS

Modeling, isotopic and geochemical data, and indirect biological indicators suggest major historical changes in atmospheric composition, particularly during the late Paleozoic but also in the late Mesozoic and Tertiary. Periods of atmospheric hyperoxia can be correlationally linked with the late Paleozoic origin of flight in insects and with contemporaneous arthropod gigantism and can be more tentatively associated with the origin of flight in three vertebrate lineages. Causal

association between periods of hyperoxia and the appearance of flight resides at two levels: biomechanical enhancement of aerodynamic force production through increases in air density and the Re , and physiological facilitation of the oxygen flux required to sustain high rates of flight metabolism. In and of themselves, increased oxygen concentrations did not act directly on particular phenotypes or behaviors. Instead, a new selective background of slowly increasing oxygen concentrations provided the opportunity (but not necessity) for the evolution and progressive enhancement of a novel locomotor capacity (26).

Falsification of such historical hypotheses is necessarily challenging, but neontological investigations of animal flight performance provide a useful perspective on paleobiological reconstructions. Postulated air densities and oxygen concentrations of ancient atmospheres can be mimicked experimentally using different combinations of oxygen and nonreactive gases (194). Similar gas mixtures can be used to investigate the effects of atmospheric hyperoxia on the biomechanics and physiology of flight in extant volant forms. Phenotypic plasticity in the respiratory system of insects (195–197) and vertebrates (198–204) suggests widespread occurrence of ontogenetic as well as short-term and chronic physiological flexibility in response to variable oxygen availability. Over evolutionary time scales, artificial selection for enhanced flight performance in *Drosophila* (205) can be imposed in different gas mixtures that decouple the effects of hyperoxia from those associated with a hyperdense flight medium (25, 194).

In modern ecological contexts, variation in altitude provides the closest analogue to historic variation in atmospheric composition. Covariance in air density and oxygen partial pressure characterizes altitudinal gradients, as would similarly characterize atmospheres with varying oxygen content and a constant quantity of nitrogen. Modern volant taxa display remarkable flight abilities under both hypoxic and hypodense conditions (61, 206–209). In natural montane contexts, wing length of birds and insects varies directly with altitude, suggesting morphological compensation for the greater power expenditure during flight at lower air densities (194, 210–213). The wide range of extant adaptations to variation in air density and oxygen availability suggests substantial evolutionary capacity to respond to variable atmospheric composition. Although considerable uncertainty still surrounds paleoatmospheric reconstructions, further study of animal flight performance in simulated ancient atmospheres is likely to prove rewarding.

ACKNOWLEDGMENTS

I thank Doug Altshuler, Bob Berner, Carl Gans, and Tim Lenton for useful comments, and acknowledge grants from the National Science Foundation for research support.

Visit the Annual Reviews home page at www.AnnualReviews.org.

LITERATURE CITED

1. Hudson JD. 1989. Palaeoatmospheres in the Phanerozoic. *J. Geol. Soc. London* 146:155–60
2. Lambert IB, Donnelly TH. 1991. Atmospheric oxygen levels in the Precambrian: a review of isotopic and geological evidence. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 97:83–91
3. Des Marais DJ, Strauss H, Summons RE, Hayes, JM. 1992. Carbon isotope evidence for the stepwise oxidation of the Proterozoic environment. *Nature* 359: 605–9
4. Knoll AH. 1992. The early evolution of eukaryotes: a geological perspective. *Science* 256:622–27
5. Canfield DE, Teske A. 1996. Late Proterozoic rise in atmospheric oxygen concentration inferred from phylogenetic and sulphur-isotope studies. *Nature* 382:127–32
6. Gilbert DL. 1996. Evolutionary aspects of atmospheric oxygen and organisms. In *Environmental Physiology*, ed. MJ Fregly, CM Blatteis, 2:1059–94. New York: Oxford Univ. Press
7. Nursall JR. 1959. Oxygen as a prerequisite to the origin of Metazoa. *Nature* 183:1170–72
8. Berkner LV, Marshall LC. 1965. On the origin and rise of oxygen concentration in the Earth's atmosphere. *J. Atmos. Sci.* 22:225–61
9. Berkner LV, Marshall LC. 1967. The rise of oxygen in the earth's atmosphere with notes on the martian atmosphere. *Adv. Geophys.* 12:309–31
10. Knoll AH, Hayes JM, Kaufman AJ, Swett K, Lambert, IB. 1986. Secular variation in carbon isotope ratios from Upper Proterozoic successions of Svalbard and East Greenland. *Nature* 321:832–38
11. Conway Morris S. 1993. The fossil record and the early evolution of the Metazoa. *Nature* 361:219–25
12. Kasting JF, Holland HD, Kump LR. 1992. Atmospheric evolution: the rise of oxygen. In *The Proterozoic Biosphere. A Multidisciplinary Study*, ed. JW Schopf, C Klein, pp. 159–63. New York: Cambridge Univ. Press
13. Kasting JF. 1993. Earth's early atmosphere. *Science* 259:920–26
14. Logan GA, Hayes JM, Hieshima GB, Summons RE. 1995. Terminal Proterozoic reorganization of biogeochemical cycles. *Nature* 376:53–56
15. Holland HD. 1984. *The Chemical Evolution of the Atmosphere and Oceans*. Princeton, NJ: Princeton Univ. Press
16. Budyko MI, Ronov AB, Yanshin AL. 1985. *History of the Earth's Atmosphere*. Berlin: Springer-Verlag
17. Schlesinger WH. 1991. *Biogeochemistry: An Analysis of Global Change*. San Diego: Academic
18. Deleted in proof
19. Tappan H. 1974. Molecular oxygen and evolution. In *Molecular Oxygen in Biology*, ed. O. Hayaishi, pp. 81–135. Amsterdam: North Holland
20. Robinson JM. 1991. Phanerozoic atmospheric reconstructions: a terrestrial perspective. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 97:51–62
21. Berner RA, Canfield DE. 1989. A new model for atmospheric oxygen over Phanerozoic time. *Am. J. Sci.* 289:333–61
22. Berner RA. 1998. The carbon cycle and CO₂ over Phanerozoic time: the role of land plants. *Philos. Trans. R. Soc. London Ser. B* 353:75–82
23. Graham JB, Dudley R, Aguilar N, Gans C. 1995. Implications of the late Palaeozoic oxygen pulse for physiology and evolution. *Nature* 375:117–20

24. Graham JB, Aguilar N, Dudley R, Gans C. 1997. The late Paleozoic atmosphere and the ecological and evolutionary physiology of tetrapods. In *Amniote Origins: Completing the Transition to Land*, ed. SS Sumida, KLM Martin, pp. 141–67. New York: Academic
25. Dudley R. 1998. Atmospheric oxygen, giant Paleozoic insects and the evolution of aerial locomotor performance. *J. Exp. Biol.* 201:1043–50
26. Gans C, Dudley R, Aguilar NM, Graham JB. 1999. Late Paleozoic atmospheres and biotic evolution. *Hist. Biol.* 13:199–219
27. Berner RA. 1990. Atmospheric carbon dioxide levels over Phanerozoic time. *Science* 249:1382–85
28. Berner RA. 1994. GEOCARB II: a revised model of atmospheric CO₂ over Phanerozoic time. *Am. J. Sci.* 294:56–91
29. Berner RA. 1997. The rise of plants and their effect on weathering and atmospheric CO₂. *Science* 276:544–46
30. Mackenzie FT, Morse JW. 1992. Sedimentary carbonates through Phanerozoic time. *Geochim. Cosmochim. Acta* 56:3281–95
31. Visscher H, Brinkhuis H, Dilcher DL, Elsik WC, Eshet Y, et al. 1996. The terminal Paleozoic fungal event: evidence of terrestrial ecosystem destabilization and collapse. *Proc. Natl. Acad. Sci. USA* 93:2155–58
32. Algeo TJ, Scheckler SE. 1998. Terrestrial-marine teleconnections in the Devonian: links between the evolution of land plants, weathering processes, and marine anoxic events. *Philos. Trans. R. Soc. London Ser. B* 353:113–30
33. Van Cappellen P, Ingall ED. 1996. Redox stabilization of the atmosphere and oceans by phosphorus-limited marine productivity. *Science* 271:493–96
34. Lenton TM, Watson AJ. 1999. Redfield revisited: 2. What regulates the oxygen content of the atmosphere? *Global Biogeochem. Cycles*. In press
35. Robinson JM. 1990. Lignin, land plants, and fungi: biological evolution affecting Phanerozoic oxygen balance. *Geology* 15:607–10
36. Watson A, Lovelock JE, Margulis L. 1978. Methanogenesis, fires and the regulation of atmospheric oxygen. *BioSystems* 10:293–98
37. Kump LR. 1989. Chemical stability of the atmosphere and ocean. *Global Planet. Change* 1:12 3–26
38. Jones TP, Chaloner WG. 1991. Fossil charcoal, its recognition and palaeoatmospheric significance. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 97:39–50
39. Rasbash DJ, Langford B. 1968. Burning of wood in atmospheres of reduced oxygen concentrations. *Combust. Flame* 12: 33–40
40. Cope MJ, Chaloner WG. 1980. Fossil charcoal as evidence of past atmospheric composition. *Nature* 283:647–49
41. Clark FRS, Russell DA. 1981. Fossil charcoal and the palaeoatmosphere. *Nature* 290:428
42. Chaloner WG. 1989. Fossil charcoal as an indicator of palaeoatmospheric oxygen level. *J. Geol. Soc. London* 146: 171–74
43. Berner RA. 1989. Drying, O₂ and mass extinction. *Nature* 340:603–4
44. Holser WT, Schönlaub HP, Attrep M Jr, Boeckelmann K, Klein P, et al. 1989. A unique geochemical record at the Permian/Triassic boundary. *Nature* 337: 39–44
45. Małkowski K, Gruszczyński M, Hoffman A, Halas S. 1989. Oceanic stable isotope composition and a scenario for the Permo-Triassic crisis. *Hist. Biol.* 2:289–309
46. Hallam A. 1991. Why was there a delayed radiation after the end-Palaeozoic extinctions? *Hist. Biol.* 5:257–62
47. Wignall PB, Hallam A. 1992. Anoxia as a cause of the Permian/Triassic mass extinction: facies evidence from northern Italy and the western United States.

- Palaeogeogr. Palaeoclimatol. Palaeoecol.* 93:21–46
48. Erwin DH. 1993. *The Great Paleozoic Crisis: Life and Death in the Permian*. New York: Columbia Univ. Press
49. Wignall PB, Hallam A. 1993. Griesbachian (Earliest Triassic) palaeoenvironmental changes in the Salt Range, Pakistan and southeast China and their bearing on the Permo-Triassic mass extinction. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 102:215–37
50. Wignall PB, Twitchett RJ. 1996. Oceanic anoxia and the End Permian mass extinction. *Science* 272:1155–58
51. Isozaki Y. 1997. Permo-Triassic boundary superanoxia and stratified superocean: records from lost deep sea. *Science* 276:235–38
52. Deleted in proof
53. Berner RA. 1991. A model for atmospheric CO₂ over Phanerozoic time. *Am. J. Sci.* 291:339–76
54. Mora CI, Driese SG, Colarusso LA. 1996. Middle to late Paleozoic atmospheric CO₂ levels from soil carbonate and organic matter. *Science* 271:1105–7
55. Retallack GJ. 1997. Early forest soils and their role in Devonian global change. *Science* 276:583–85
56. McElwain JC. 1998. Do fossil plants signal palaeoatmospheric CO₂ concentration in the geological past? *Philos. Trans. R. Soc. London Ser. B* 353:83–96
57. Beerling DJ, Woodward FI, Lomas MR, Wills MA, Quick WP, Valdes PJ. 1998. The influence of Carboniferous palaeo-atmospheres on plant function: an experimental and modelling assessment. *Philos. Trans. R. Soc. London Ser. B* 353:131–40
58. Edwards D. 1998. Climate signals in Palaeozoic land plants. *Philos. Trans. R. Soc. London Ser. B* 353:141–57
59. Vogel S. 1994. *Life in Moving Fluids: The Physical Biology of Flow*. Princeton, NJ: Princeton Univ. Press
60. Norberg UM. 1990. *Vertebrate Flight*. Berlin: Springer-Verlag
61. Dudley R. 2000. *The Biomechanics of Insect Flight: Form, Function, Evolution*. Princeton, NJ: Princeton Univ. Press
62. Whalley P, Jarzembowski EA. 1981. A new assessment of *Rhyniella*, the earliest known insect, from the Devonian of Rhynie, Scotland. *Nature* 291:317
63. Shear WA, Grierson JD, Rolfe WDI, Smith EL, Norton RA. 1984. Early land animals in North America: evidence from Devonian age arthropods from Gilboa, New York. *Science* 224:492–94
64. Labandeira CC, Beall BS, Hueber FM. 1988. Early insect diversification: evidence from a Lower Devonian bristletail from Québec. *Science* 242:913–16
65. Jeram AJ, Selden PA, Edwards D. 1990. Land animals in the Silurian: arachnids and myriapods from Shropshire, England. *Science* 250:658–61
66. Nelson CR, Tidwell WD. 1987. *Brodioptera stricklani* n.sp. (Megasecoptera: Brodiopteridae), a new fossil insect from the Upper Manning Canyon Shale Formation, Utah (lowermost Namurian B). *Psyche* 94:309–16
67. Brauckmann C, Zessin W. 1989. Neue Meganeuridae aus dem Namurian von Hagen-Vorhalle (BRD) und die Phylogenie der Meganisoptera. *Dtsch. Entomol. Z. N.F.* 36:177–215
68. Wootton RJ. 1981. Palaeozoic insects. *Annu. Rev. Entomol.* 26:319–44
69. Wootton RJ. 1990. Major insect radiations. In *Major Evolutionary Radiations*, ed. PD Taylor, GP Larwood, Systematics Assoc. Special Volume 42:187–208. Oxford, UK: Clarendon
70. Kukalová-Peck J. 1991. Fossil history and the evolution of hexapod structures. In *The Insects of Australia*, ed. CSIRO, 1:141–79. Ithaca, NY: Cornell Univ. Press. 2nd ed.
71. Labandeira CC, Sepkoski JJ. 1993. Insect diversity in the fossil record. *Science* 261:310–15

72. Boudreaux BH. 1979. *Arthropod Phylogeny with Special Reference to Insects*. New York: Wiley & Sons
73. Hennig W. 1981. *Insect Phylogeny*. Chichester, UK: Wiley & Sons
74. Kristensen NP. 1981. Phylogeny of insect orders. *Annu. Rev. Entomol.* 26:135–57
75. Kristensen NP. 1989. Insect phylogeny based on morphological evidence. In *The Hierarchy of Life*, ed. B Fernholm, K Bremer, H. Jörnvall, pp. 295–306. Amsterdam: Elsevier Sci.
76. Kristensen NP. 1991. Phylogeny of extant hexapods. See Ref. 70, pp. 125–40
77. Kristensen NP. 1997. The groundplan and basal diversification of the hexapods. In *Arthropod Relationships*, ed. RA For- tey, RH Thomas, pp. 281–93. London: Chapman & Hall
78. Brodsky AK. 1994. *The Evolution of Insect Flight*. Oxford, UK: Oxford Univ. Press
79. Crampton GC. 1916. The phylogenetic origin and the nature of the wings of insects according to the paranotal theory. *J. NY Entomol. Soc.* 24:1–39
80. Forbes WTM. 1943. The origin of wings and venational types in insects. *Am. Midl. Nat.* 29:381–405
81. Hamilton KGA. 1971. The insect wing, Part I. Origin and development of wings from notal lobes. *J. Kans. Entomol. Soc.* 44:421–33
82. Rasnitsyn AP. 1981. A modified paranotal theory of insect wing origin. *J. Morphol.* 168:331–38
83. Quartau JA. 1985. On some objections to the paranotal theory on the origin of insect wings. *Bol. Soc. Port. Entomol. Suppl.* 1:359–71
84. Quartau JA. 1986. An overview of the paranotal theory on the origin of the insect wings. *Publ. Inst. Zool. Dr. Augusto Nobre, Fac. Cienc. Porto* 194:1–42
85. Wootton RJ. 1986. The origin of insect flight: Where are we now? *Antenna* 10:82–86
86. Bitsch J. 1994. The morphological groundplan of Hexapoda: critical review of recent concepts. *Ann. Soc. Entomol. Fr.* 30:103–29
87. Woodworth CW. 1906. The wing veins of insects. *Univ. Calif. Bull. Entomol.* 1:1–52
88. Wigglesworth VB. 1973. Evolution of insect wings and flight. *Nature* 246:127–29
89. Wigglesworth VB. 1976. The evolution of insect flight. In *Insect Flight*, ed. RC Rainey, pp. 255–69. Oxford, UK: Blackwell
90. Kukalová-Peck J. 1978. Origin and evolution of insect wings and their relation to metamorphosis, as documented by the fossil record. *J. Morphol.* 156:53–126
91. Kukalová-Peck J. 1983. Origin of the insect wing and wing articulation from the arthropodan leg. *Can. J. Zool.* 61:1618–69
92. Kukalová-Peck J. 1992. The “Uniramia” do not exist: the ground plan of the Pterygota as revealed by Permian Diaphanopteroidea from Russia (Insecta: Paleodictyopteroidea). *Can. J. Zool.* 70: 236–55
93. Kukalová-Peck J. 1997. Arthropod phylogeny and ‘basal’ morphological structures. See Ref. 77, pp. 249–68
94. Kukalová-Peck J. 1997. Mazon Creek insect fossils: the origin of insect wings and clues about the origin of insect metamorphosis. In *Richardson’s Guide to the Fossil Fauna of Mazon Creek*, ed. CW Shabic, AA Hay, pp. 194–207. Chicago: Northeastern Illinois Univ. Press
95. Averof M, Cohen, SM. 1997. Evolutionary origin of insect wings from ancestral gills. *Nature* 385:627–30
96. Little C. 1983. *The Colonisation of Land. Origins and Adaptations of Terrestrial Animals*. Cambridge, UK: Cambridge Univ. Press
97. Resh VH, Solem JO. 1984. Phylogenetic

- relationships and evolutionary adaptations of aquatic insects. In *An Introduction to the Aquatic Insects of North America*, ed. RW Merritt, KW Cummins, pp. 66–75. Dubuque, IA: Kendall/Hunt. 2nd ed.
98. Messner B. 1988. Sind die Insekten primäre oder sekundäre Wasserbewohner? *Dtsch. Entomol. Z. N.F.* 35:355–60
99. Pritchard G, McKee MH, Pike EM, Scrimgeour GJ, Zloty J. 1993. Did the first insects live in water or in air? *Biol. J. Linn. Soc.* 49:31–44
100. Hinton HE. 1968. Spiracular gills. *Adv. Insect Physiol.* 5:65–161
101. Carpenter FM. 1966. The Lower Permian insects of Kansas. Part 11. The orders Protorthoptera and Orthoptera. *Psyche* 73:46–88
102. Kukalová J. 1968. Permian mayfly nymphs. *Psyche* 75:310–27
103. Carroll SB, Weatherbee SD, Langeland JA. 1995. Homeotic genes and the regulation and evolution of insect wing number. *Nature* 375:58–61
104. Becker EG. 1952. The problem of the origin and development of the wing in insects. Chapter 1. Precursors of the insect wing. *Vestn. Mosk. Univ. Ser. Phys. Math. Nat. Sci.* 9:59–68
105. Wilson DM. 1962. Bifunctional muscles in the thorax of grasshoppers. *J. Exp. Biol.* 39:669–77
106. Ewer DW. 1963. On insect flight. *J. Entomol. Soc. S. Afr.* 26:3–13
107. Ewer DW, Nayler LS. 1967. The pterothoracic musculature of *Deropeltis erythrocephala*, a cockroach with a wingless female, and the origin of wing movements in insects. *J. Entomol. Soc. S. Afr.* 30:18–33
108. Fournier CR, Randall JB. 1982. Studies on cockroach flight: the role of continuous neural activation of non-flight muscles. *J. Exp. Zool.* 221:143–54
109. Wigglesworth VB. 1963. The origin of flight in insects. *Proc. R. Entomol. Soc. London* 28:23–32
110. Alexander RD, Brown WL. 1963. Mating behavior and the origin of insect wings. *Occas. Pap. Mus. Zool. Univ. Mich.* 628:1–19
111. Alexander RD. 1964. The evolution of mating behaviour in arthropods. *Symp. R. Entomol. Soc. London* 2:78–94
112. Whalley PS. 1979. New species of Protorthoptera and Protodonata (Insecta) from the Upper Carboniferous of Britain, with a comment on the origin of wings. *Bull. Br. Mus. Nat. Hist.* 32:85–90
113. Douglas MM. 1981. Thermoregulatory significance of thoracic lobes in the evolution of insect wings. *Science* 211:84–86
114. Kingsolver JG, Koehl MAR. 1985. Aerodynamics, thermoregulation, and the evolution of insect wings: differential scaling and evolutionary change. *Evolution* 39:488–504
115. Kingsolver JG, Koehl MAR. 1994. Selective factors in the evolution of insect wings. *Annu. Rev. Entomol.* 39:425–51
116. Bradley, JC. 1942. The origin and significance of metamorphosis and wings among insects. In *Proc. 8th Am. Sci. Congr.* pp. 303–9. Washington DC: Dept. State
117. Mamayev BM. 1977. The gravitational hypothesis of the origin of insects. *Entomol. Rev.* 54:13–17
118. Marden JH, Kramer MG. 1994. Surface-skimming stoneflies: a possible intermediate stage in insect flight evolution. *Science* 266:427–30
119. Marden JH, Kramer MG. 1995. Locomotor performance of insects with rudimentary wings. *Nature* 377:332–34
120. Thomas ALR, Norberg RÅ. 1996. Skimming the surface—the origin of flight in insects? *Trends Ecol. Evol.* 11:187–88
121. Samways MJ. 1994. ‘Sailing’ on the water surface by adult male *Enallagma nigridorsum* Selys (Zygoptera: Coenagrionidae). *Odonatologica* 23:175–78

122. Kramer MG, Marden JH. 1997. Almost airborne. *Nature* 385:403–4
123. Will KW. 1995. Plecopteran surface-skimming and insect flight evolution. *Science* 270:1684–85
124. Ruffieux L, Elouard J-M, Sartori M. 1998. Flightlessness in mayflies and its relevance to hypotheses on the origin of insect flight. *Proc. R. Soc. London Ser. B* 265:2135–40
125. Flower JW. 1964. On the origin of flight in insects. *J. Insect Physiol.* 10:81–88
126. Wootton RJ. 1976. The fossil record and insect flight. See Ref. 89, pp. 235–54
127. Ritzmann RE, Fourtner CR. 1980. Flight activity initiated via giant interneurons of the cockroach: evidence for bifunctional trigger interneurons. *Science* 210:443–45
128. Ritzmann RE. 1984. The cockroach escape response. In *Neural Mechanisms of Startle Behavior*, ed. RC Eaton, pp. 93–131. New York: Plenum
129. Edwards JS, Reddy GR. 1986. Mechanosensory appendages in the firebrat (*Thermobia domestica*, Thysanura): a prototype system for terrestrial predator invasion. *J. Comp. Neurol.* 243:535–46
130. Bristowe WS. 1958. *The World of Spiders*. London: Collins
131. Edwards JS. 1985. Predator evasion and the origin of insect flight: an exercise in evolutionary neuroethology. *Soc. Neurosci. Abstr.* 11:497
132. Edwards JS, Palka J. 1991. Insect neural evolution—a fugue or an opera? *Semin. Neurosci.* 3:391–98
133. Edwards JS. 1992. Giant interneurons and the origin of insect flight. In *Nervous Systems: Principles of Design and Function*, ed. RN Singh, pp. 485–95. Bombay: Wiley Eastern
134. Edwards JS. 1997. The evolution of insect flight: implications for the evolution of the nervous system. *Brain Behav. Evol.* 50:8–12
135. Ganihar D, Libersat F, Wendler G, Camhi, JM. 1994. Wind-evoked evasive responses in flying cockroaches. *J. Comp. Physiol. A* 175:49–65
136. Kutsch W, Breidbach O. 1994. Homologous structures in the nervous systems of Arthropoda. *Adv. Insect Physiol.* 24:1–11
137. Libersat F. 1994. The dorsal giant interneurons mediate evasive behavior in flying cockroaches. *J. Exp. Biol.* 197:405–11
138. Rolfe WD. 1980. Early invertebrate terrestrial fossils. In *The Terrestrial Environment and the Origin of Land Vertebrates*, ed. AL Panchen, Systematics Assoc. Special Vol. No. 15, pp. 117–57. London: Academic
139. Rolfe WD. 1985. Early terrestrial arthropods: a fragmentary record. *Philos. Trans. R. Soc. London Ser. B* 309:207–18
140. Carroll RL. 1988. *Vertebrate Paleontology and Evolution*. New York: Freeman
141. Shear WA, Kukalová-Peck J. 1990. The ecology of Paleozoic terrestrial arthropods: the fossil evidence. *Can. J. Zool.* 68:1807–34
142. Behrensmeyer AK, Damuth JD, DiMichele WA, Sues H-D, Wing SL. 1992. *Terrestrial Ecosystems through Time: Evolutionary Paleocology of Terrestrial Plants and Animals*. Chicago: Univ. Chicago Press
143. Carpenter FM. 1971. Adaptations among Paleozoic insects. *Proc. N. Am. Paleontol. Conv.* 1969:1236–51
144. Burnham L. 1983. Studies on Upper Carboniferous insects: 1. The Geraridae (Order Protorthoptera). *Psyche* 90:1–57
145. Kevan PG, Chaloner WG, Savile DBO. 1975. Interrelationships of early terrestrial arthropods and plants. *Paleontology* 18:391–417
146. Chaloner WG, Sheerin A. 1979. Devonian macrofloras. *Spec. Pap. Palaeontol.* 23:145–61
147. Gensel PG, Andrews HN. 1984. *Plant Life in the Devonian*. New York: Praeger
148. Stewart WN. 1983. *Paleobotany and the Evolution of Plants*. Cambridge, UK: Cambridge Univ. Press
149. Raven JA. 1986. Evolution of plant life forms. In *On the Economy of Plant Form*

- and Function*, ed. TJ Givnish, pp. 421–92. Cambridge, UK: Cambridge Univ. Press
150. Kenrick P, Crane PR. 1997. The origin and early evolution of plants on land. *Nature* 389:33–39
151. Strong DR, Lawton JH, Southwood R. 1984. *Insects on Plants*. Oxford, UK: Blackwell
152. Rasnitsyn AP, Krassilov VA. 1996. First find of pollen grains in the gut of Permian insects. *Paleontol. J.* 30:484–90
153. Labandeira CC, Phillips TL. 1996. A Carboniferous insect gall: insight into early ecologic history of the Holometabola. *Proc. Natl. Acad. Sci. USA* 93:8470–74
154. Labandeira CC, Phillips TL. 1996. Insect fluid-feeding on Upper Pennsylvanian tree ferns (Palaedictyoptera, Marattiales) and the early history of the piercing-and-sucking functional feeding group. *Ann. Entomol. Soc. Am.* 89:157–83
155. Ellington CP. 1991. Aerodynamics and the origin of insect flight. *Adv. Insect Physiol.* 23:171–210
156. Ennos AR. 1989. The effect of size on the optimal shapes of gliding insects and seeds. *J. Zool. London* 219:61–69
157. Haupt H. 1941. Die ältesten geflügelten Insekten und ihre Beziehungen zur Fauna der Jetztzeit. *Z. Naturwiss.* (Halle) 94:60–121
158. Smart J. 1971. Palaeoecological factors affecting the origin of winged insects. *Proc. 13th Int. Congr. Entomol.* 1:304–6
159. Harlé É, Harlé A. 1911. Le vol de grands reptiles et insectes disparus semble indiquer une pression atmosphérique élevée. *Bull. Soc. Geol. Fr. Ser. 4* 11:18–21
160. Kammer AE, Heinrich B. 1978. Insect flight metabolism. *Adv. Insect Physiol.* 13:133–228
161. Casey TM. 1989. Oxygen consumption during flight. In *Insect Flight*, ed. GJ Goldsworthy, CH Wheeler, pp. 257–72. Boca Raton, FL: CRC Press
162. Weis-Fogh T. 1964. Diffusion in insect wing muscle, the most active tissue known. *J. Exp. Biol.* 41:229–56
163. Weis-Fogh T. 1964. Functional design of the tracheal system of flying insects as compared with the avian lung. *J. Exp. Biol.* 41:207–27
164. Kestler P. 1984. Respiration and respiratory water loss. In *Environmental Physiology and Biochemistry of Insects*, ed. KH Hoffman, pp. 137–83. Berlin: Springer-Verlag
165. Mill PJ. 1985. Structure and physiology of the respiratory system. In *Comprehensive Insect Physiology, Biochemistry, and Pharmacology*, ed. GA Kerkut, LI Gilbert, 3:517–93. Oxford, UK: Pergamon
166. Harrison JF, Lighton JRB. 1998. Oxygen-sensitive flight metabolism in the dragonfly *Erythemis simplicicollis*. *J. Exp. Biol.* 201:1739–44
167. Kukalová-Peck J. 1985. Ephemeroïd wing venation based upon new gigantic Carboniferous mayflies and basic morphology, phylogeny, and metamorphosis of pterygote insects (Insecta, Ephemeroïda). *Can. J. Zool.* 63:933–55
168. Carpenter FM. 1992. *Treatise on Invertebrate Paleontology. Part R, Arthropoda 4, Volumes 3 and 4 (Hexapoda)*. Lawrence, KS: Univ. Kansas Press
169. Kukalová-Peck J. 1987. New Carboniferous Diplura, Monura, and Thysanura, the hexapod ground plan, and the role of thoracic lobes in the origin of wings (Insecta). *Can. J. Zool.* 65:2327–45
170. Rolfe WDI. 1969. Arthropleurida. In *Treatise on Invertebrate Paleontology. Arthropoda 4: Crustacea (Except Ostracoda): Myriapoda*, ed. RC Moore, C Teichert, pp. 607–20. Lawrence, KS: Univ. Kansas Press
171. Kraus O. 1974. On the morphology of Paleozoic diplopods. *Symp. Zool. Soc. London* 32:13–22
172. Briggs, DEG. 1985. Gigantism in Palaeozoic arthropods. *Spec. Pap. Paleontol.* 33:157

173. Vermeij GJ. 1987. *Evolution and Escalation*. Princeton, NJ: Princeton Univ. Press
174. Ruten MG. 1966. Geologic data on atmospheric history. *Paleogeogr. Palaeoclimatol. Palaeoecol.* 2:47–57
175. Schidlowski M. 1971. Probleme der atmosphärischen Evolution im Präkambrium. *Geol. Rundsch.* 60:1351–84
176. Jepsen GL. 1970. Bat origins and evolution. In *Biology of Bats*, ed. WA Wimsatt, 1:1–64. London: Academic
177. Altringham JD. 1996. *Bats: Biology and Behaviour*. Oxford, UK: Oxford Univ. Press
178. Chiappe LM. 1995. The first 85 million years of avian evolution. *Nature* 378:349–55
179. Qiang J, Currie PJ, Norell MA, Shu-An J. 1998. Two feathered dinosaurs from northeastern China. *Nature* 393:753–61
180. Padian K, Chiappe LM. 1998. The origin and early evolution of birds. *Biol. Rev.* 73:1–42
181. Wild R. 1984. Flugsaurier aus der Obertrias von Italien. *Naturwissenschaften* 71:1–11
182. Wellnhofer P. 1991. *The Illustrated Encyclopedia of Pterosaurs*. London: Salamander
183. Ruben J. 1991. Reptilian physiology and the flight capacity of *Archaeopteryx*. *Evolution* 45:1–17
184. Ruben J. 1993. Powered flight in *Archaeopteryx*: response to Speakman. *Evolution* 47:935–38
185. Speakman JR. 1993. Flight capabilities in *Archaeopteryx*. *Evolution* 41:336–40
186. Norberg UM. 1985. Evolution of vertebrate flight: an aerodynamic model for the transition from gliding to active flight. *Am. Nat.* 126:303–27
187. Rayner JMV. 1988. The evolution of vertebrate flight. *Biol. J. Linn. Soc.* 34:269–87
188. Bennett SC. 1997. The arboreal leaping theory of the origin of pterosaur flight. *Hist. Biol.* 12:265–90
189. Eaton RC, ed. 1984. *Neural Mechanisms of Startle Behavior*. New York: Plenum
190. Tuttle MD, Stevenson D. 1982. Growth and survival of bats. In *Ecology of Bats*, ed. TH Kunz, pp. 105–50. New York: Plenum
191. Pomeroy D. 1990. Why fly? The possible benefits for lower mortality. *Biol. J. Linn. Soc.* 40:53–65
192. Rose MR. 1991. *Evolutionary Biology of Aging*. New York: Oxford Univ. Press
193. Holmes DJ, Austad SN. 1995. The evolution of avian senescence patterns: implications for understanding primary aging processes. *Am. Zool.* 35:307–17
194. Dudley R, Chai P. 1996. Animal flight mechanics in physically variable gas mixtures. *J. Exp. Biol.* 199:1881–85
195. Loudon C. 1988. Development of *Tenebrio molitor* in low oxygen levels. *J. Insect Physiol.* 34:97–103
196. Loudon C. 1989. Tracheal hypertrophy in mealworms: design and plasticity in oxygen supply systems. *J. Exp. Biol.* 147:217–35
197. Greenberg S, Ar A. 1996. Effects of chronic hypoxia, normoxia and hyperoxia on larval development in the beetle *Tenebrio molitor*. *J. Insect Physiol.* 42:991–96
198. Temple GF, Metcalfe J. 1970. The effects of increased incubator oxygen tension on capillary development in the chick chorioallantois. *Condor* 90:187–92
199. McGrath JJ. 1971. Acclimation response of pigeons to simulated high altitude. *J. Appl. Physiol.* 31:274–76
200. Pionetti J-M, Bouverot P. 1977. Effects of acclimation to altitude on oxygen affinity and organic phosphate concentrations in pigeon blood. *Life Sci.* 20:1207–12
201. Metcalfe J, McCutcheon, IE, Francisco DL, Metzner AB, Welch JE. 1981. Oxygen availability and growth of the chick embryo. *Respir. Physiol.* 46:81–88
202. Williams JB, Swift K. 1988. Oxygen consumption and growth of Northern

- Bobwhite embryos under normoxic and hyperoxic conditions. *Condor* 90:187–92
203. Bigard AX, Brunet A, Guezennec C-Y, Monod H. 1991. Effects of chronic hypoxia and endurance training on muscle capillarity in rats. *Pflügers Arch.* 419: 225–29
204. Hoppeler H, Desplanches D. 1992. Muscle structural modifications in hypoxia. *Int. J. Sports Med.* 13(Suppl. 1):S166–68
205. Weber KE. 1996. Large genetic change at small fitness cost in large populations of *Drosophila melanogaster* selected for wind tunnel flight: rethinking fitness surfaces. *Genetics* 144:205–13
206. Galun R, Fraenkel G. 1961. The effect of low atmospheric pressure on adult *Aedes aegyptii* and on housefly pupae. *J. Insect Physiol.* 7:161–76
207. Dudley R. 1995. Extraordinary flight performance of orchid bees (Apidae: Euglossini) hovering in heliox (80% He/20% O₂). *J. Exp. Biol.* 198:1065–70
208. Chai P, Dudley R. 1995. Limits to vertebrate locomotor energetics suggested by hummingbirds hovering in heliox. *Nature* 377:722–25
209. Chai P, Dudley R. 1996. Limits to flight energetics of hummingbirds hovering in hypodense and hypoxic gas mixtures. *J. Exp. Biol.* 199:2285–95
210. Stalker HD, Carson HL. 1948. An altitudinal transect of *Drosophila robusta* Sturtevant. *Evolution* 2:295–305
211. Hamilton TH. 1961. The adaptive significances of intraspecific trends of variation in wing length and body size among bird species. *Evolution* 15:180–95
212. Feinsinger P, Colwell RK, Terborgh J, Chaplin SB. 1979. Elevation and the morphology, flight energetics, and foraging ecology of tropical hummingbirds. *Am. Nat.* 113:481–97
213. Epting RJ. 1980. Functional dependence of the power for hovering on wing disc loading in hummingbirds. *Physiol. Zool.* 53:347–57

Copyright of Annual Review of Physiology is the property of Annual Reviews Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.