

Higher categories of Tettigoniidea (katydids*) represented in FSCA

Names used are those recognized by [OSF Online](#).

T. J. Walker and Katherine Arguez (August 2017)

Order Orthoptera, Suborder Ensifera, Infraorder Tettigoniidea

Superfamily Hagloidea

Family Prophalangopsidae (1 genus, 3 spp.)

Superfamily Rhabdophoroidea

Family Rhabdophoridae

Subfamily Aemodogryllinae (1 genus, 1 sp.)

Subfamily Ceuthophilinae (4 genera, 8 spp.)

Subfamily Gammarotettiginae (1 genus, 1 sp.)

Subfamily Tropidischinae (1 genus, 1 sp.)

Superfamily Stenopelmatoidea

Family Anostomatidae (1 genus)

Family Gryllacrididae (1 genus, 1 sp.)

Family Stenopelmatidae (1 genus, 1 sp.)

Superfamily Tettigoniioidea

Family Tettigoniidae

Subfamily Conocephalinae

Tribe Conocephalini (3 genera, 42 spp.)

Tribe Copiphorini (13 genera, 44 spp.)

Subfamily Hetrodinae (2 genera)

Subfamily Listroscelidinae (1 genus, 2 spp.)

Subfamily Meconematinae (2 genera, 2 spp.)

Subfamily Pterochrozinae (1 genus, 2 spp.)

Subfamily Saginae (1 genus, 1 sp.)

Subfamily Tettigoniinae (27 genera, 74 spp.)

Family Phaneropteridae**

Subfamily Mecopodinae (no identified holdings)

Subfamily Phaneropterinae (26 genera, 85 spp.)

Subfamily Phyllophorinae (no identified holdings)

Subfamily Pseudophyllinae (21 genera, 26 spp.)

* There is no colloquial name that covers only the insect taxa on this sheet. "*Katy-did*" is the rendering of the loud, persistent calling of one species of Pseudophyllinae and a common name for that species. However, in North America, *katydid* is often used more broadly as a "common" name for the superfamily Tettigoniioidea. Here the concept is stretched further to apply to all taxa in the Infraorder Tettigoniidea.

** The boundaries among the subfamilies of Phaneropteridae are unclear, in part because two of the main subfamilies (Mecopodinae and Phyllophorinae) are found almost exclusively in the Old World.