



Bulletin of the
**Oregon
Entomological
Society**

Steiroxys strepens (Noisy Shieldback) at Woodruff Meadows

text and photos by Ron Lyons

Bulletin of the Oregon Entomological Society, 2019(3), pp. 1–4.

Steiroxys strepens (Noisy Shieldback) at Woodruff Meadows

text and photos by Ron Lyons

Steiroxys is a genus of shield-backed katydids (Orthoptera: Tettigoniidae: Tettigoniinae) restricted to western North America and the southern portion of western Canada. Only 4 species have been described so far:

Steiroxys borealis Scudder, 1894,
Steiroxys pallidipalpus (Thomas, 1872),
Steiroxys strepens Fulton 1930, and
Steiroxys trilineata (Thomas, 1870).

In their revision of the shield-backed katydids, Rentz and Birchim (1968: 144) write, “Members of this diurnal genus have always been baffling to systematists at the specific level. . . . Only study of large series will show the true relationships. . . .” There is evidence that all 4 species, at least as currently defined, can be found in Oregon. The genus is in serious need of revision and additional species are being processed.

Steiroxys strepens (Figure 1–6) was described by Fulton (1930: 627–630) using specimens collected from two locations in Oregon: Jackson’s Hill (6 miles north of Corvallis) in Benton County and Woodruff Meadows in Jackson County. Jackson’s Hill is now called Dimple Hill (see Hietala-Henschell [2017]). Woodruff Meadows is west of Highway 62 just south of the boundary between Jackson and Douglas Counties, a few miles northwest of Prospect. Fulton did not indicate that individuals had been found anywhere else.

Steiroxys strepens is a diurnal katydid with short flight wings—like many other species of shield-backed katydids, it cannot fly. The type specimens were from Jackson’s Hill as were 5 male and 7 female paratypes. Eleven additional male paratypes were collected at Woodruff Meadows. The Jackson’s Hill specimens used in the description were collected July 12, 1922. (Fulton mentions that first and second instar nymphs were found the following year on April 26.) The date that the Woodruff Meadows specimens were collected was not specifically mentioned under the description but was probably August 5, 1922, mentioned later in a discussion of that area (p. 636). It is interesting that Fulton does not mention

having collected any females from Woodruff Meadows. All of his specimens were green, except for one male collected from Woodruff Meadows which appeared to have been light brown. Although Fulton worked at the Oregon Agricultural College (now Oregon State University) at the time, none of his *Steiroxys* specimens are present in the Oregon State Arthropod Collection (the paper was published after Fulton took up a position at North Carolina State College). Photographs of the male type and female allotype can be found on the Orthoptera Species File website, <<http://orthoptera.speciesfile.org/Common/basic/Taxa.aspx?TaxonNameID=1142430>>.



Figure 1: Light brown male of *Steiroxys strepens* photographed at Woodruff Meadows on August 22, 2019.



Figure 2. Left: Dark brown male *Steiroxys strepens* photographed August 22, 2019. Above: Cerci of the same male.

Males of *Steiroxys strepens* can be separated from those of other species of *Steiroxys* by the shape of their cerci. Fulton (p. 628) writes “Cerci gently curved throughout on outer border; **terminal portion finger-shaped, nearly straight and with rounded apex** [emphasis added]. A curved spine projects perpendicularly inward just beyond the middle of the cercus and terminates in a sharp claw.” He also writes that “The male cerci of the Woodruff Meadows series have on the average a stouter and shorter apical portion and a longer and more slender inner spine.”

In 2019, I made four visits to the east side of Woodruff Meadows (August 22, September 5, September 12, and September 26) finding *Steiroxys strepens* on each occasion—the number on each

date, based on my photographs, were 10, 3, 1 and 1 respectively; I only found males, 5 of them, on the first trip. I found green and brown individuals—the brown individuals ranged from light to dark brown; only one female had a noticeable dorsal stripe (Figure 6). The appearance of the male cerci (Figures 2 and 3) and the female sub-genital plate (Figure 7) agree in general with the descriptions and accompanying figures in Fulton’s work. Fulton described the call of *Steiroxys strepens*; I was not aware of any calls on my trips (field start times ranged from 10:40 to 13:30).

In the Oregon State Arthropod Collection there is a male collected at Union Creek on VII 27 '45 by H.A. Scullen (OSAC 693637; Figure 8) (this location is just east of Woodruff



Figure 3. Left: Green male photographed on August 22, 2019. Above: Cerci of the same male.



Figure 4. Green female photographed September 5, 2019.



Figure 5. Brown female photographed September 12, 2019.

Meadows). There are also 7 females collected from Mary's Peak near Corvallis. The features of the male and the female specimens are consistent with Fulton's work.

Rentz and Birchim (p. 144) say "Great variation in the shape of the cerci occurs even among individuals from a single locality." From Fulton's work and my limited sample, I do not believe this to be the case for *Steiroxys strepens*. For that reason, I have excluded these 2 reports from Oregon that would have expanded its range in the state:

- 1) *Steiroxys strepens* has been reported from the H.J. Andrews Forest near Blue River in Lane County (Lightfoot 1981).
- 2) on bugnet, a male *Steiroxys* photographed in Union County has been identified as *Steiroxys strepens* (<<https://bugguide.net/node/view/702410/bgimage>>).

In both cases, **based solely on the material presented**, the cerci do not appear to be consistent with Fulton's description. (David Lightfoot recently confirmed [priv. comm. October 2019] that the species from the Andrews Forest is not *Steiroxys strepens*.) Outside of Oregon, Miskelly (2012) reported that a *Steiroxys* that looks very much like *Steiroxys strepens* occurs on Vancouver Island in British Columbia.

There are at least 2 shield-backed katydid known from, or adjacent to, the area where *Steiroxys strepens* is currently known in Oregon. *Tessellana tessellata*, a non-native diurnal shield-backed katydid, has expanded northward since its introduction into central California around 1950, and occurs well into Washington (Lyons 2012, Walker 2019), and now perhaps even southern British Columbia. This katydid is widespread in western Oregon and has distinctive flight wings of variable length. In Woodruff



Figure 6. This female *Steiroxys strepens* has the most extreme color pattern of any of the individuals I photographed. It is the only one with a dorsal stripe, noticeable mainly because of contrast between the pale dorsum of the pronotum and its dark sides.



Figure 7. Sub-genital plate of green female found September 26, 2019. The image was taken through glass and is slightly overexposed. Part of the ovipositor can be seen at the bottom of the image.

Meadows, it is more common than *Steiroxys strepens* but is smaller and easily overlooked with all the grasshopper activity. The second species, *Idiostatus gurneyi* described in Rentz (1973: 42–48), is a diurnal species that has been recorded at Crater Lake, west of Woodruff Meadows. Rentz points out that *Idiostatus gurneyi* “shares many characters with species of the genus *Steiroxys*.” In particular, the male cerci (Figure 8) look very much like those described for *Steiroxys strepens*. Currently, it is “known from montane peaks from northern California and southern Oregon.”

At Woodruff Meadows, I found at least 7 of the grasshopper species found by Fulton (p. 637): *Chloealtis aspasma*, *Oedaleonotus pacificus*, *Melanoplus bivittatus* (Two-striped Grasshopper), *Melanoplus validus*, *Camnula pellucida* (Clear-winged Grasshopper), *Cratypedes neglectus* (Pronotal Range Grasshopper), and *Trimerotropis fontana*. In addition, I have a couple pictures that may be *Melanoplus lovetti*, a species described by Fulton in the same paper and, so far, known only from Woodruff Meadows. I also found some other grasshopper species not specifically mentioned by Fulton.

Fulton (1930: 636–637) discussed the landscape of Woodruff Meadows. He refers to a “. . . striking zonation of the vegetation. The center contained some surface water and was covered with a high sedge and large lupines. Surrounding this was a wide zone of damp ground with shorter swamp grasses, and another zone of dryer ground bordering the forest and covered mostly by herbs.” I am not sure how big the area was that Fulton described. The central area, which I did not get to, has a confluence of streams according to my map. Most of the rest is damp to dry mixed woodland with scattered open areas and several large fenced off meadows, although the fences I came across are in disrepair. The evidence of free-roaming cattle was apparent everywhere I went, but I only saw them once as a herd moved through one of the large meadows.

Fulton pointed out that there are many small natural prairies on the west slopes of the Cascades. He indicated, “Their isolation and diverse conditions as to moisture and temperature tend to give them a more or less individual fauna,” making them worthy of

further investigation.

References

- Fulton, B.B. 1930. Notes on Oregon Orthoptera, with Descriptions of New Species and Races. *Annals of the Entomological Society of America* 23(4): 611–641.
- Hietala-Henschell, K. 2017. Species Fact Sheet (for *Chloealtis aspasma* Rehn and Hebard 1919). Xerces Society: 16 p. (This is an updated version of the fact sheet prepared by G. Brenner dated August 2006.) (Download the WORD file from <<https://www.fs.fed.us/rb/sfpnw/issssp/documents5/sfs-iiior-chloealtis-aspasma-2017-09.doc>>.)
- Lightfoot, D.C. 1986. Invertebrates of the H.J. Andrews Experimental Forest, Western Cascades, Oregon: III. The Orthoptera (Grasshoppers and Crickets). USDA Forest Service Pacific Northwest Research Station Research Note PNW-443: 23 p. (Download PDF from <https://www.fs.fed.us/pnw/pubs/pnw_rn443.pdf>.)
- Lyons, R. 2012. *Tessellana tessellata* in Oregon (Orthoptera: Tettigoniidae). *Bulletin of the Oregon Entomological Society* Fall 2012: 5–6.
- Miskelly, J.W. 2012. Updated Checklist of the Orthoptera of British Columbia. *Journal of the Entomological Society of British Columbia* 109: 24–30. (Download the PDF from <<https://journal.entsocbc.ca/index.php/journal/article/download/231/370>>.)
- Rentz, D.C. 1973. The Shield-backed Katydid of the Genus *Idiostatus*. *Memoirs of the American Entomological Society* 29: 211 p.
- Rentz, D.C. and J.D. Birchim. 1968. Revisionary Studies in the Nearctic Decticinae. *Memoirs of the Pacific Coast Entomological Society* 3: 173 p.
- Walker, T.J. 2019. Singing Insects of North America (SINA) [<<http://entnemdept.ifas.ufl.edu/walker/Buzz/>>]: *Tessellana tessellata* (<<http://entnemdept.ufl.edu/walker/buzz/316a.htm>> [accessed September 2019].)



Figure 8. Left: Cerci of male *Steiroxys strepens* collected at Union Creek in Jackson County on VII 27 '45 by H.A. Scullen (OSAC 693637). Right: Cerci of male *Idiostatus gurneyi* photographed on Mount Ashland September 5, 2012.

Corrections Added to Reprint

Two errors were noticed up in the preparation of this reprint.

1. The text on page 3, paragraph 2, item 2 reads:

2) on bugnet ...

The text should refer to the website as bugguide not bugnet.

2. On page 4, paragraph 1 the text reads:

Idiostatus gurneyi ... recorded at Crater Lake, west of
Woodruff Meadows.

Crater Lake [National Park] is east of Woodruff Meadows.

Ron Lyons
October 15, 2020